

**STEGLATRO® ▼**  
(ertugliflozin)

**PRESCRIBING INFORMATION**

Refer to Summary of Product Characteristics (SmPC) before prescribing.

**Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in Google Play or Apple App Store. Adverse events should also be reported to MSD, UK (Tel: 01992 467272). By clicking the above link you will leave the MSD website and be taken to the MHRA website.**

**PRESENTATION**

Film-coated tablets containing 5 mg or 15 mg of ertugliflozin (as ertugliflozin L-pyroglytamic acid)

**USES**

Type 2 diabetes mellitus as an adjunct to diet and exercise to improve glycaemic control:

- as monotherapy in patients for whom the use of metformin is considered inappropriate due to intolerance or contraindications.
- with other anti-diabetic medicinal products.

**DOSAGE AND ADMINISTRATION**

Recommended starting dose is 5 mg once daily. Increase to 15 mg once daily if necessary. Assess renal function and correct volume depletion prior to initiation.

**Renal impairment:** Do not initiate in patients with eGFR <60ml/min/1.73m<sup>2</sup>. Discontinue when eGFR persistently <45 ml/min/1.73 m<sup>2</sup>.

**Hepatic impairment:** mild or moderate impairment: no dose adjustment required; severe impairment: not recommended. **Elderly:** ≥ 65 years: no dose adjustment required; ≥75 years: limited data. **Children** <18 years: no data.

**CONTRA-INDICATIONS**

Hypersensitivity to active substance or excipients.

**PRECAUTIONS**

Do not use in patients with type 1 diabetes. Symptomatic hypotension may occur upon initiation of therapy, particularly in patients with impaired renal function. Assess volume status prior to initiation. Consider temporary interruption of therapy if volume depletion occurs. Use with caution in patients at risk of DKA. If DKA is suspected or diagnosed, discontinue ertugliflozin. Interrupt therapy in patients hospitalised for major surgical procedures or acute serious medical illness until stabilised. A small increase in the risk of lower limb amputation (primarily of the toe) has been reported. Counsel all patients on routine preventative foot care and maintaining adequate hydration. Consider stopping treatment should lower-extremity skin ulcers, osteomyelitis or gangrene develop. Assess renal function prior to initiation of therapy and periodically thereafter. Discontinue ertugliflozin if eGFR falls persistently <45ml/min/1.73m<sup>2</sup>. Monitor patients for

genital mycotic infections and treat appropriately. Consider temporary interruption of ertugliflozin when treating pyelonephritis or urosepsis. Fournier's gangrene (FG) has been reported with SGLT2 inhibitors in both sexes and may be preceded by urogenital infection or perineal abscess. Advise patients to seek medical attention in the event of fever or malaise with pain, tenderness, erythema or swelling in the genital or perineal area. If FG is suspected, discontinue ertugliflozin and treat promptly. No experience in NYHA class III-IV. Contains lactose monohydrate. Do not use in patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption. Do not use urine glucose or 1,5-AG assay to monitor glycaemic control.

**Drug interactions:** Diuretics, insulin, sulphonyureas.

**Pregnancy and lactation:** Not recommended.

**SIDE EFFECTS**

**Refer to SmPC for complete information on side-effects.**

**Very common (≥ 1/10):** vulvovaginal mycotic infection and other female genital mycotic infections.

**Common (≥ 1/100 to < 1/10):** Balanitis candida and other male genital mycotic infections, hypoglycaemia, volume depletion, increased urination, vulvovaginal pruritus, thirst, serum lipid changes, increased haemoglobin and BUN.

**Uncommon (≥ 1/1,000 to <1/100):** Dysuria, blood creatinine increased, glomerular filtration decreased.

**Rare (≥1/10,000 to <1/1,000):** DKA.

**Not known:** Fournier's gangrene.

**PACKAGE QUANTITIES AND BASIC NHS COST**

5 mg x 28: £29.40

15 mg x 28: £29.40

**Marketing Authorisation numbers**

5 mg x 28: EU/1/18/1267/002

15 mg x 28: EU/1/18/1267/008

**Marketing Authorisation Holder**

Merck Sharp & Dohme B.V.  
Waarderweg 39  
2031 BN Haarlem  
The Netherlands

**Legal category: POM**

**Date of review of prescribing information: June 2019**